

Data Sheet

CellGenix® Recombinant Human Interleukin-3 (rh IL-3) Preclinical Grade - Order No.: 1402-010 (10 µg), 1402-050 (50 µg)

Product Characteristics

Source	<i>E. coli</i>
Description	Human Interleukin-3, accession # P08700, Ala20-Phe152 N-terminal Met Molecular mass 15.2 kDa
Formulation	Lyophilized from a 0.2 µm-filtered solution containing 10 mM sodium phosphate and 50 mM sodium chloride, pH 7.2.
Intended use	For preclinical <i>ex vivo</i> use. Not intended for therapeutic use.

Quality Parameters

Activity	≥ 0.9 x 10 ⁶ IU/mg calibrated against NIBSC #91/510 Measured in a cell proliferation assay using an IL-3-dependent cell line, TF-1
Purity	≥ 95 %, as determined by SDS-PAGE (under reducing and non-reducing conditions, visualized by Coomassie staining) and RP-HPLC
Endotoxin	< 1000 EU/mg, as determined by LAL gel clot test
Sterility	Sterility test of the vial product (direct inoculation)
Mass per vial	1402-010: 10 µg, 1402-050: 50 µg
Animal-derived component-free	ADCF Level 2: The final product contains neither animal- nor human-derived materials. ADCF Level 2 cytokines are produced in our dedicated animal-free facility. No animal-derived components are used throughout the complete production process. All ADCF Level 2 cytokines are produced in <i>E. coli</i> .

Shipment & Storage

Transport	Ambient temperature. Please refer to Technote (www.cellgenix.com)
Expiry	3 years from date of shipment
Storage & Stability	Store lyophilized cytokine at -20°C to -80 °C. Store a 100 µg/ml reconstituted cytokine solution: <ul style="list-style-type: none">• 4 weeks at 2°C to 8°C under sterile conditions after reconstitution. Store in the original container.• 4 months at -20°C to -80°C under sterile conditions after reconstitution. Store in 80 µl aliquots in polypropylene cryogenic vials. Avoid repeated freeze/thaw cycles.

Handling Instructions

Reconstitution	Recommended in sterile water to a final concentration of 100 µg/ml (for 10 µg vials) or 250 µg/ml (for 50 µg vials).
Dilution	Recommended in CellGenix® serum-free media. For dilution with protein free medium, a carrier protein (0.1–1 % albumin or 1–10 % appropriate serum) has to be included. Failure to dilute product according to these instructions may result in loss of activity.

Quality Statement

Final manufacturing steps and QC are performed in a GMP facility.